

Breaking the Silence: A Case Study on Bulimia Nervosa in a Young Adult Male

Adeena Zakariya,
Student, Psychology Department,
Dr Sunaina Kaur Maan
Assistant Professor,
The Bhopal School of Social Sciences, Bhopal, MP, India

Abstract

Bulimia nervosa is a severe and potentially life-threatening eating disorder characterized by recurrent episodes of binge eating followed by compensatory behaviors to prevent weight gain. This case study examines the presentation, diagnosis, treatment, and outcomes of a 23-year-old male medical student diagnosed with bulimia nervosa. The patient reported a one-year history of binge eating, self-induced vomiting, and excessive exercise, driven by body dissatisfaction and low self-esteem. He also had a history of substance abuse and a prior diagnosis of bipolar disorder. A comprehensive assessment revealed psychological comorbidities, including anxiety and depressive symptoms, alongside physical complications such as electrolyte imbalances.

The treatment approach included a multidisciplinary plan encompassing cognitive-behavioral therapy (CBT), pharmacological interventions, and family support. Over six months, the patient demonstrated gradual improvements in eating patterns, self-perception, and mental health. This case highlights the complexity of bulimia nervosa as a multifactorial condition requiring integrated care strategies. It also underscores the importance of early intervention, individualized treatment, and continued research to optimize outcomes and reduce the stigma associated with eating disorders.

Keywords: Bulimia nervosa, eating disorder, substance abuse, bipolar disorder, cognitive-behavioral therapy

Introduction

Bulimia nervosa is a complex and serious eating disorder characterized by recurrent episodes of binge eating followed by compensatory behaviors such as self-induced vomiting, excessive exercise, or misuse of laxatives and diuretics (American Psychiatric Association [APA], 2013). Individuals with bulimia nervosa often experience profound guilt, shame, and a perceived loss of control over their eating habits, perpetuating the cycle of disordered behaviors.

Recognized as a distinct psychological condition in the late 20th century, bulimia nervosa is influenced by a combination of biological, psychological, and sociocultural factors. It is often associated with distorted body image, low self-esteem, and a preoccupation with weight and appearance. While the condition can affect individuals across all demographics, it is most commonly diagnosed in adolescents and young adults, with a higher prevalence among females (Smink et al., 2012).

If untreated, bulimia nervosa can lead to significant medical complications, including electrolyte imbalances, gastrointestinal disturbances, severe dental erosion, and an increased risk of comorbid psychiatric disorders such as anxiety, depression, and substance use (Mehler & Brown, 2015). A comprehensive approach to treatment, including psychotherapy, medical monitoring, nutritional counseling, and pharmacological interventions, is critical for recovery. This case study explores the diagnosis, treatment, and outcomes of a 23-year-old male patient with bulimia nervosa, highlighting the importance of early intervention and individualized care.

Case Presentation

Subject Details:

- **Name:** Zain Javadd
- **Age:** 23 years
- **Gender:** Male
- **Professional Status:** Medical student
- **Diagnosed Problem:** Bulimia nervosa

Background:

The patient, a 23-year-old medical student from Bhopal, comes from a well-educated and supportive family. His father is a doctor, and his mother manages the household. He is the second eldest of four siblings, with an elder sister who is an MBBS doctor, a twin brother also studying medicine, and a younger sister in high school. The patient described a generally positive relationship with his siblings but reported occasional conflicts with his parents.

The patient's struggles with body image began during his weight loss journey, where he lost 38 kilograms (from 108 kg to 70 kg) in three months through unhealthy eating habits. This

rapid weight loss led to the development of bulimia nervosa, characterized by binge eating followed by self-induced vomiting. The patient also had a history of substance abuse and a prior diagnosis of bipolar disorder.

Symptoms and Diagnosis

The patient exhibited the following symptoms:

1. **Binge Eating:** Consuming large amounts of food in a short period, followed by feelings of guilt and shame.
2. **Purging Behaviors:** Self-induced vomiting and excessive exercise to prevent weight gain.
3. **Physical Signs:** Calluses on knuckles, bloodshot eyes, and significant weight fluctuations.
4. **Psychological Symptoms:** Low self-esteem, distorted body image, and mood swings.

The patient was diagnosed with bulimia nervosa based on the DSM-5 criteria (APA, 2013). His history of bipolar disorder and substance abuse added complexity to the diagnosis, necessitating a comprehensive treatment approach.

Treatment Plan

The treatment plan included the following components:

1. **Cognitive-Behavioral Therapy (CBT):** Focused on identifying and challenging distorted thoughts about body image and eating behaviors. The patient learned healthier coping mechanisms and strategies to break the binge-purge cycle.
2. **Pharmacological Interventions:**
 - **Prozac (fluoxetine):** An SSRI to reduce binge-purge frequency and alleviate depressive symptoms.
 - **Zoloft (sertraline):** Another SSRI to stabilize mood and manage anxiety.
 - **Tricyclic Antidepressants (TCAs):** Used to address severe depressive episodes and enhance appetite regulation.
3. **Family Involvement:** Psychoeducation sessions were conducted to help the family understand the disorder and provide a supportive environment.

4. **Nutritional Counseling:** Guidance on establishing balanced and sustainable eating habits.

Progress and Outcomes

Over six months, the patient demonstrated significant improvements:

- Reduced frequency of binge-purge episodes.
- Improved mood stability and self-esteem.
- Healthier eating patterns and weight management.
- Increased engagement in academic and social activities.

Despite these improvements, the patient continued to experience occasional struggles with body image and self-esteem, highlighting the need for long-term follow-up and support.

Discussion

This case study illustrates the multifactorial nature of bulimia nervosa, emphasizing the interplay of psychological, physiological, and environmental factors. The patient's history of rapid weight loss, substance abuse, and bipolar disorder underscores the complexity of the disorder and the importance of a comprehensive, individualized treatment approach.

The integration of CBT, pharmacological interventions, and family support proved effective in addressing the core symptoms of bulimia nervosa and associated comorbidities. Early diagnosis and intervention were critical in mitigating the physical and psychological effects of the disorder.

Conclusion

Bulimia nervosa is a challenging and multifaceted condition that requires a compassionate, evidence-based approach to treatment. This case study highlights the importance of early intervention, individualized care, and long-term support in achieving sustained recovery. By promoting awareness and reducing stigma, clinicians and caregivers can encourage individuals with eating disorders to seek timely help and improve their quality of life.

References

American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Arlington, VA: American Psychiatric Publishing.

Mehler, P. S., & Brown, C. (2015). Anorexia nervosa – medical complications. *Journal of Eating Disorders*, 3(1), 11. <https://doi.org/10.1186/s40337-015-0040-8>

Smink, F. R., van Hoeken, D., & Hoek, H. W. (2012). Epidemiology of eating disorders: Incidence, prevalence, and mortality rates. *Current Psychiatry Reports*, 14(4), 406-414. <https://doi.org/10.1007/s11920-012-0282-y>