

Manifestation of Anxiety and Depression in Individuals

Pradeep Kumar Kushwah

Student

Santripty Gautam

Student

Dr Sunaina Kaur Maan

Assistant Professor

The Bhopal School of Social Sciences, Bhopal, MP, India

Abstract

The exact cause of Generalized Anxiety Disorder (GAD) remains incompletely understood. Current hypotheses suggest that fluctuations in neurotransmitter levels, including serotonin, dopamine, and norepinephrine, contribute to alterations in patients' psychological states. This case study examines the manifestations of anxiety and depression in Nishay, a 22-year-old male from a rural background residing in Bhopal for work. Nishay exhibits symptoms of anxiety and depression that have persisted for several years, alongside multiple health conditions. As the eldest child in his family, he bears significant familial responsibilities, contributing to stress, anxiety, loss of interest in activities, poor appetite, sleep disturbances, anger issues, and occasional self-harm.

Despite practicing mindfulness, breathing exercises, and Jacobson's Progressive Muscle Relaxation (JPMR), as well as engaging in yoga and physical exercise, Nishay continues to experience debilitating symptoms that disrupt his social, personal, and professional life. Diagnostic tools, including the Generalized Anxiety Disorder-7 (GAD-7), Beck Depression Inventory (BDI), Self-Injury Questionnaire (SIQ), and Eating Disorder Examination Questionnaire (EDE-Q), were used to assess the severity of his conditions. The results confirmed significant levels of anxiety, depression, eating disorders, and self-destructive behaviors.

This study underscores that anxiety and depression are not merely states of mind that can be alleviated through physical exercise or positive thinking alone. Instead, they involve a complex interplay of psychological and physiological factors, including chemical imbalances in the brain, which can lead to persistent and debilitating symptoms. By exploring Nishay's case in detail, this study highlights the broader implications of these conditions and emphasizes the need for comprehensive treatment approaches. The findings suggest that while physical therapy

alone cannot cure GAD or depression, it may help alleviate some physical symptoms associated with these conditions.

Keywords: Anxiety, Depression, Stress, Sleep Disturbance, Self-harm, Eating Disorders, Physical Exercise

Introduction

Anxiety disorders and depression are prevalent mental health conditions that significantly impact individuals' daily lives and overall well-being. Anxiety is characterized by excessive worry, fear, and physical symptoms, while depression is marked by persistent sadness, loss of interest in activities, and various physical manifestations. These conditions often coexist, creating a complex interplay of symptoms that can severely impair an individual's quality of life.

Generalized Anxiety Disorder (GAD) is defined by excessive fear and worry, whereas depression involves persistent low mood and a lack of interest in previously enjoyable activities. Both conditions can lead to secondary health issues, such as sleep disturbances, appetite changes, and self-harming behaviors. The complexities of these disorders extend beyond their psychological effects, profoundly influencing physical health and daily functioning.

This case study focuses on Nishay, a 22-year-old male from a rural background, who exhibits severe symptoms of anxiety and depression. Over the past few years, Nishay has faced numerous health challenges, compounded by his role as the eldest son in his family. This position has placed significant financial and emotional stress on him, exacerbating his symptoms. His persistent anxiety has led to disruptions in his social, personal, and professional life. Loss of interest in activities, constant isolation, and difficulty managing anger have further complicated his condition. Nishay's introverted nature and limited social circle have also made him a target of bullying, while his need for control has resulted in restrictive eating patterns and significant weight loss.

The Complex Interplay of Factors Contributing to Nishay's Condition

1. Family Responsibilities and Socio-Economic Stress:

- As the eldest son, Nishay faces immense pressure to support his family, exacerbating his stress and anxiety.
- Financial instability adds another layer of stress, creating a cycle of worry that is difficult to break.

2. Health Conditions and Physical Symptoms:

- Nishay's ongoing health issues contribute to his anxiety, as chronic illness often leads to increased worry about health and financial stability.
- Physical symptoms of anxiety, such as poor appetite and sleep disturbances, have directly impacted his overall well-being.

3. Mental Health Challenges:

- The co-occurrence of anxiety and depression in Nishay's case highlights the complexity of his condition. Symptoms such as loss of interest in activities and self-harming behaviors add to the burden of anxiety.
- Anger issues and irritability further indicate high levels of distress and frustration.

Despite Nishay's efforts to manage his symptoms through yoga, physical exercise, and self-help methods, his condition has not significantly improved. This underscores the fact that anxiety and depression are not merely states of mind that can be resolved through physical activity or positive thinking. Instead, they involve a complex interplay of psychological and physiological factors, including chemical imbalances in the brain, which can result in persistent and debilitating symptoms.

To assess the severity of Nishay's conditions, several diagnostic tools were administered, including the Generalized Anxiety Disorder-7 (GAD-7), Beck Depression Inventory (BDI), Self-Injury Questionnaire (SIQ), and Eating Disorder Examination Questionnaire (EDE-Q). The results confirmed the severity of his anxiety, depression, eating disorder, and self-destructive behaviors. This case study highlights the intricate nature of these conditions and the necessity of comprehensive treatment approaches that go beyond surface-level interventions.

Methodology

This section outlines the steps taken to conduct the case study on anxiety and depression, focusing on Nishay, a 22-year-old male. It details the participant selection process, data

collection methods, and diagnostic tools used to assess the severity of symptoms and gather comprehensive information about the subject's condition.

Participants:

The primary participant in this case study is Nishay, who exhibits traits of severe anxiety and depression. Nishay was selected based on observable symptoms and his willingness to participate in the study. His background, including health conditions, familial responsibilities, and socio-economic status, provided a rich context for understanding the complexities of his condition.

Data Collection:

Data collection involved observational methods to ensure a thorough understanding of Nishay's condition.

- **Observation:** Observational sessions were conducted in various settings, including Nishay's home and workplace, to monitor his behavior and interactions. These sessions aimed to identify environmental factors contributing to his anxiety and depression in real-time.

Diagnostic Tools:

The following tools were used to assess the severity of Nishay's symptoms:

1. **Generalized Anxiety Disorder-7 (GAD-7):** A self-report questionnaire designed to screen for and measure the severity of generalized anxiety disorder.
2. **Beck Depression Inventory (BDI):** A self-report questionnaire that measures the severity and behavioral manifestations of depression.
3. **Self-Injury Questionnaire (SIQ):** A 54-item self-report questionnaire assessing self-injurious behaviors.
4. **Eating Disorder Examination Questionnaire (EDE-Q):** A self-report instrument measuring eating disorder behaviors and attitudes.

Analysis

The analysis of Nishay's case involved examining qualitative and quantitative data to identify common themes, differences, and correlations related to his anxiety and depression.

Common Themes:

1. **Persistent Anxiety Symptoms:** Nishay has experienced anxiety symptoms for several years, including excessive worry, palpitations, sweating, sleep disturbances, and difficulties in social and work settings.
2. **Impact of Socio-Economic Stressors:** Financial instability and familial responsibilities significantly contribute to Nishay's anxiety, leading to persistent worry, low mood, and self-destructive behaviors.
3. **Ineffectiveness of Self-Help Methods:** Despite efforts to manage symptoms through yoga and physical exercise, Nishay's condition has not significantly improved.
4. **Chemical Imbalance:** The persistence and severity of Nishay's symptoms suggest a potential link to chemical imbalances in the brain, highlighting the need for medication as part of his treatment plan.

Differences:

1. **Rural Background and Limited Awareness:** Nishay's rural background and limited understanding of mental health have influenced his reliance on traditional and self-help methods.
2. **Health Conditions:** Nishay's multiple health issues add complexity to his condition, impacting his ability to engage in coping mechanisms effectively.
3. **Personal Motivation:** Despite challenges, Nishay demonstrates a strong motivation to manage his symptoms, reflecting a proactive approach to self-improvement.

Correlations:

1. **Anxiety and Financial Stress:** A clear correlation exists between Nishay's financial stress and the severity of his anxiety symptoms.
2. **Sleep Disturbances and Worry:** Sleep disturbances are closely linked to Nishay's excessive worry, creating a cycle that impacts his overall well-being.

3. **Social and Work Life Impact:** Nishay's symptoms significantly affect his social and professional life, reinforcing the need for comprehensive treatment.

Conclusion

Nishay's case highlights the multidimensional nature of anxiety disorders, depression, and associated conditions, emphasizing their profound impact on an individual's life. Despite his efforts to manage symptoms through self-help methods, Nishay continues to experience moderate to severe anxiety and depression, underscoring the need for a comprehensive treatment approach.

This study emphasizes that anxiety and depression are not merely states of mind that can be resolved through physical exercise or positive thinking. Instead, they involve a complex interplay of psychological, physiological, and socio-economic factors. The persistence of Nishay's symptoms suggests that chemical imbalances in the brain play a critical role, necessitating the inclusion of medication in his treatment plan.

Furthermore, Nishay's rural background and limited awareness of mental health issues highlight the importance of increasing access to professional mental health services in underdeveloped areas. Education and awareness programs can help individuals like Nishay better understand their conditions and seek appropriate treatment.

References

- Kessler, R. C., Chiu, W. T., Demler, O., & Walters, E. E. (2005). Prevalence, severity, and comorbidity of twelve-month DSM-IV disorders in the National Comorbidity Survey Replication (NCS-R). *Archives of General Psychiatry*, 62(6), 617-627.
- Spitzer, R. L., Kroenke, K., Williams, J. B., & Löwe, B. (2006). A brief measure for assessing generalized anxiety disorder: the GAD-7. *Archives of Internal Medicine*, 166(10), 1092-1097.

Appendices

Appendix A: GAD-7 Questionnaire

Instructions: Fill in the actual responses from Nishay.

Questions:

1. Feeling nervous, anxious, or on edge. – Nearly every day
2. Not being able to stop or control worrying. – Nearly every day
3. Worrying too much about different things. – Nearly every day
4. Trouble relaxing. – Nearly every day
5. Being so restless that it is hard to sit still. – Nearly every day
6. Becoming easily annoyed or irritable. – Nearly every day
7. Feeling afraid as if something awful might happen. – Nearly every day